

Dyadic Measures of the Parent–Child Relationship During the Transition to Adolescence and Glycemic Control in Children With Type 1 Diabetes

BARBARA J. ANDERSON, PHD
GRAYSON HOLMBECK, PHD
RONALD J. IANNOTTI, PHD
SIRIPOOM V. MCKAY, MD
AMANDA LOCHRIE, PHD
LISA K. VOLKENING, MA
LORI LAFFEL, MD

To identify aspects of family behavior associated with glycemic control in youth with type 1 diabetes mellitus during the transition to adolescence, the authors studied 121 9- to 14-year-olds ($M = 12.1$ yrs) and their parents, who completed the Diabetes Family Conflict Scale (DFCS) and the Diabetes Family Responsibility Questionnaire (DFRQ). From the DFRQ, the authors derived 2 dyadic variables, frequency of agreement (exact parent and child concurrence

about who was responsible for a task) and frequency of discordance (opposite parent and child reports about responsibility). The authors divided the cohort into Younger ($n = 57$, $M = 10.6$ yrs) and Older ($n = 64$, $M = 13.5$ yrs) groups. Family conflict was significantly related to glycemic control in the entire cohort and in both the Younger and Older groups. However, only in the Younger group was Agreement related to glycemic control, with higher Agreement associated with

Barbara J. Anderson, PhD, and Siripoom V. McKay, MD, Texas Children's Hospital; Grayson Holmbeck, PhD, Loyola University Chicago; Ronald J. Iannotti, PhD, National Institute of Child Health and Human Development, Bethesda, Maryland; Amanda Lochrie, PhD, Nemours Children's Clinic, Jacksonville, Florida; Lisa K. Volkening, MA, and Lori Laffel, MD, Joslin Diabetes Center, Boston, Massachusetts.

The authors represent the Steering Committee of the Family Management of Childhood Diabetes Study.

This research was supported by the intramural research program of the National Institutes of Health, National Institute of Child Health and Human Development. The following investigators and institutions made up the steering committee of the Family Management of Childhood Diabetes multisite trial: Jill Weissberg-Benchell, PhD, and Grayson Holmbeck, PhD (Contract N01-HD-4-3363), Children's Memorial

Hospital, Chicago; Bruce Simons-Morton, EdD, Tonja R. Nansel, PhD, Ronald J. Iannotti, PhD, and Rusan Chen, PhD, National Institute of Child Health and Human Development, Bethesda, Maryland; Barbara Anderson, PhD (Contract N01-HD-4-3362), Texas Children's Hospital, Houston, Texas; Tim Wysocki, PhD, and Amanda Lochrie, PhD (Contract N01-HD-4-3361), Nemours Children's Clinic, Jacksonville, Florida; Lori Laffel, MD, MPH, Deborah Butler, MSW, Korey Hood, PhD, and Lisa Volkening, MA (Contract N01-HD-4-3364), Joslin Diabetes Center, Boston; and Cheryl McDonnell, PhD, and MaryAnn D'Elia (Contract N01-HD-3-3360), James Bell Associates, Arlington, Virginia.

Correspondence concerning this article should be addressed to Barbara J. Anderson, PhD, Baylor College of Medicine, Texas Children's Hospital, 1100 Bates Street, Houston, TX 77030. E-mail: bja@bcm.tmc.edu

better glycemic control. Findings suggest that Agreement about sharing of diabetes responsibilities may be an important target for family-based interventions aiming to optimize glycemic control in preteen youth.

Keywords: adolescence, type 1 diabetes, glycemic control, dyadic family variables, conflict

Type 1 diabetes mellitus (T1DM), the second most common chronic illness of childhood after asthma, has a complex medical management regimen that requires multiple daily injections of insulin, pricking a finger several times a day to monitor blood glucose levels, and adjusting insulin dosages on the basis of the child's blood glucose level, food intake, and physical activity (Silverstein et al., 2005). These behavioral tasks are carried out to keep blood glucose levels as close to normal as possible. Adherence to the diabetes regimen is important because keeping blood sugar levels as close to normal as possible, from as early in the disease course as possible, helps to prevent or delay the devastating long-term complications of T1DM, such as blindness, kidney disease, amputations, heart attacks, and stroke (Diabetes Control and Complications Trial Research Group, 1993; Epidemiology of Diabetes Interventions and Complications Research Group, 2001). Moreover, adhering to these regimen requirements is especially challenging when they intersect and clash with developmentally typical behaviors encountered at different stages of child development (Anderson & Brackett, 2005; Piazza-Waggoner et al., 2008).

Optimal glycemic control is the most difficult to establish and maintain during the early adolescent period because of the normal insulin resistance that occurs during puberty (Amiel, Sherwin, Simonson, Lauritano, & Tamborlane, 1986). In addition to this basic biological phenomenon, the normal developmental tasks of early adoles-

cence involving transitions in family roles and peer relationships often interfere with adherence to the diabetes treatment regimen (Anderson & Auslander, 1989; Wysocki et al., 1996). Young adolescents frequently seek a new level of separation from their parents while simultaneously intensifying attachments to peers (Levitt, Guacci-Franco, & Levitt, 1993).

Recent family studies of youth with T1DM have documented an erosion of parental involvement and support for diabetes management tasks during the early adolescent years (Anderson, Auslander, Jung, Miller, & Santiago, 1990; Anderson, Ho, Brackett, Finkelstein, & Laffel, 1997; Schilling, Knaf, & Grey, 2006; Skinner, Murphy, & Huws-Thomas, 2005; Wysocki et al., 1996). Moreover, empirical research has documented a steady decrease in adherence to diabetes treatment during early adolescence (Johnson, Silverstein, Rosenbloom, Carter, & Cunningham, 1986; Harris et al., 2000). However, empirical studies have also shown that young adolescents who have more parental involvement, monitoring, and teamwork in their diabetes management tend to achieve and maintain better diabetes outcomes (Allen, Tennen, McGrade, Affleck, & Ratzan, 1983; Anderson et al., 1997; Ingersoll, Orr, Herrold, & Golden, 1986; Palmer et al., 2004; Wiebe et al., 2005).

In the general child development literature, it is well documented that conflict between parents and children often increases during the transition to adolescence (Holmbeck, 1996). Both diabetes-specific family conflict and general family conflict have been associated with poor adherence and poor glycemic control during the early adolescent years (Anderson et al., 2002; Hauser et al., 1990). In a qualitative study of transfer of diabetes management responsibilities and conflict in parent-child dyads over the adolescent period, Schilling et al. (2006) reported that parents of 8- to 11-year-old children with diabetes took active steps to transfer responsibility for diabetes management to

their children, with only moderate amounts of conflict. Moreover, parents of 12- to 15-year-old youth continued to transfer increasing amounts of responsibility to their children despite increasing levels of conflict in the dyad.

Because of the deterioration in glycemic control, the increase in family conflict, and the shifts that occur in how parents and their older school-age children share responsibilities for the complex tasks demanded in the daily management of T1DM, behavioral interventions to enhance health outcomes in youth with T1DM have recently focused on the parent-child dyad (Anderson, Brackett, Ho, & Laffel, 1999; Ellis et al., 2005; Wysocki, Harris, Buckloh, Mertlich, et al., 2006).

Dyadic measures provide a perspective into family relationships that is not possible with self-report measures that are limited to personal behavior or with global assessments of family functioning (Cook & Kenny, 2006). In an effort to identify aspects of family behavior associated with glycemic control in youth with T1DM during the transition to adolescence, we measured diabetes-specific parent-child conflict and parent-child dyadic agreement and disagreement regarding which specific family member had primary responsibility for a range of different diabetes management tasks critical to health outcomes in young adolescents. We hypothesized that better glycemic control would be associated with lower diabetes-specific family conflict, higher dyadic agreement, and lower dyadic discordance in youth with T1DM and their parents. We also sought to explore age-related differences in how these family behaviors were associated with glycemic control in the 9- to 14-year-old youth in this study.

METHOD

Participants

Children with T1DM and their primary caregivers were recruited from four large

pediatric tertiary care diabetes centers in Boston; Chicago; Houston, Texas; and Jacksonville, Florida, to participate in a multisite pilot and feasibility study of an intervention designed to optimize family adaptation to childhood diabetes during late childhood and early adolescence. The participating parent or guardian was the adult primarily responsible for the child's diabetes management; only 1 parent participated per family. The data reported here were derived from the baseline assessment, which occurred before assignment of families to different treatment groups.

Trained research staff at each site reviewed medical record data to identify eligible patients with upcoming clinic appointments. Eligible families were then recruited both at clinic visits and by telephone. To examine age differences in diabetes-specific family interactions over the transition to early adolescence, we recruited youth between 9 and 14.5 years of age at baseline. Other youth eligibility criteria included T1DM duration of at least 1 year; insulin dose of at least 0.5 units/kg/day; at least 2 clinic visits at the diabetes center during the past year; mean hemoglobin A1c (HbA1c) of less than 13.0% during the prior 8 months; no other major chronic disease or cognitive disability; and no psychiatric hospitalization within the prior 6 months. Enrollment criteria for parents or guardians included at least a fifth-grade reading fluency in English; no diagnosis of psychosis, substance use disorder, major depression, or bipolar disorder; no psychiatric hospitalization in the prior 6 months; and working telephone service. Because the intervention to be tested in this pilot study required participants to engage in family problem-solving processes, parents and youth with depression or serious mental health diagnoses were excluded. All parents or legal guardians signed institution-approved informed consent forms, and all youth assented to participate in the study using each center's approved procedures.

A total of 167 eligible families were invited to participate in the study, and 131 (78%) initially agreed to participate. However, 10 families declined participation when contacted to schedule the baseline assessment, resulting in a final sample of 121 families (73%), with 29–31 families at each of the four clinical sites. Because this pilot and feasibility study was designed to assess the ability of the research team at each of the four sites to implement a clinic-based intervention with a representative sample of families at each site, we did not perform power analyses to establish the final sample size.

Procedure

Trained, two-person interviewing teams who were not affiliated with the clinics conducted the baseline assessments in families' homes with the parent and youth. Parents and youth completed several assessment measures simultaneously, but with different interviewers and in different rooms to allow for privacy of responses. Data from two of the psychosocial assessment instruments completed at the baseline assessment are reported here. These two measures are the Diabetes Family Responsibility Questionnaire (DFRQ; Anderson et al., 1990) and the Diabetes Family Conflict Scale (DFCS; Hood, Butler, Anderson, & Laffel, 2007). Parents and youth also both completed the following assessment measures: the Pediatric Quality of Life Inventory Generic Core Scales and Diabetes Module (Varni et al., 2003; Varni, Seid, & Kurtin, 2001); the Hypoglycemia Fear Survey—Worry Subscale (Cox, Irvine, Gonder-Frederick, Nowacek, & Butterfield, 1987); and the Blood Glucose Monitoring Communication survey (Hood, Butler, Volkening, Anderson, & Laffel, 2004). Youth also completed the Beck Depression Inventory for Children (Beck, Beck, & Jolly, 2001) or the Children's Depression Inventory (Kovacs, 1985), the Diabetes Management Self-Efficacy Scale (Iannotti et al., 2006), and several surveys assessing

youth perception of parenting style and involvement (Barber, 1996; Jackson, Henriksen, & Foshee, 1998; Nansel, Rovner, et al., 2009; Smetana & Asquith, 1994). Parents also completed the Child Maturity Scale (Hartos, Eitel, & Simons-Morton, 2001); the Confusion, Hubbub, and Order Scale (Matheny, Wachs, Ludwig, & Phillips, 1995); and the Diabetes Management Outcomes Expectations scale (Iannotti et al., 2006). For a complete description of the pilot and feasibility intervention study, see Nansel, Anderson, et al. (2009).

Measures

Diabetes Family Responsibility Questionnaire

The DFRQ is a 17-item questionnaire that measures parent involvement in diabetes management tasks (Anderson et al., 1990). For each of the 17 diabetes management tasks, respondents rated the responsibility for the task as belonging primarily to the child (1), being shared about equally between child and parent (2), or belonging primarily to the parent (3). Higher scores indicate increasing levels of parent responsibility for diabetes management. Acceptable internal consistency and test-retest reliability have previously been reported (Anderson et al., 1990). The alpha coefficients for the present sample were adequate, with an alpha of .67 for parents and an alpha of .73 for youth.

Within each family, we compared the parent and child responses to each of the 17 DFRQ items, following the method for deriving dyadic variables described by Anderson et al. (1990). This method captures extreme reports as recommended for the DFRQ and yields two derived dyadic variables: agreement and discordance. Agreement occurred when the child and parent agreed exactly in their report of who had responsibility for a specific diabetes management task (e.g., both respondents reported that the parent had primary responsibility for the task). Discordance occurred

when the child and the parent had the exact opposite report of who had responsibility for a specific diabetes management task (e.g., one person reported that the parent had primary responsibility and the other person reported that the child had primary responsibility). We summed the number of items for which there was exact parent-child agreement to obtain a total agreement score. Similarly, we summed the number of items for which there was parent-child discordance (exact opposite reports) to obtain a total discordance score. Therefore, agreement and discordance scores could range from 0 to 17.

Table 1 illustrates dyadic agreement and discordance per item on the DFRQ according to parent and child response, following the method suggested by Anderson et al. (1990). Agreement occurred when the child and parent agreed exactly in their report of who takes responsibility for a specific diabetes management task; this is represented in the three cells along the diagonal from left to right. Discordance occurred when the report of responsibility between the parent and child was completely opposite; this is represented in two cells, the upper right and lower left. A third scenario exists, in which a dyad could have partial agreement (or partial discordance) in their responses for a specific diabetes management task. For example, this would occur if one person said that the parent had

primary responsibility and the other person said that the responsibility was shared. This is represented by the remaining cells in the table (the empty cells).

Diabetes Family Conflict Scale

Youth and parents completed a 15-item version of the DFCS (Hood et al., 2007; Rubin, Young-Hyman, & Peyrot, 1989). Each parent and child indicated the amount of family conflict that occurred for each of 15 aspects of diabetes management. The score is the sum of the items on which any amount of conflict was endorsed. Alphas reported for the scale have been strong: youth = .85, parent = .81 (Hood et al., 2007). The alpha coefficients for the present sample were .90 for parents and .95 for youth.

The glycosylated hemoglobin (HbA1c) blood test estimates average glycemic concentration over the prior 2 to 3 months (Chase, 2006). The patients attended their regular diabetes clinic appointment approximately 2 weeks after the in-home baseline assessment. Patient blood samples were obtained by fingerstick at their regular clinic appointments. Samples were shipped to a central lab at the Joslin Diabetes Center (Boston) for processing using the Tosoh High Performance Liquid Chromatography 2.2 method (Tosoh Corporation, Foster City, CA). Joslin is a reference laboratory for this assay, which has been

Table 1

Derivation of Dyadic Variables of Agreement and Discordance According to Child and Parent Response for Each DFRQ Item

Child report	Parent report		
	1	2	3
1. Child	Agreement		Discordance
2. Shared		Agreement	
3. Parent	Discordance		Agreement

Note. This table depicts how the variables of agreement and discordance were derived for a parent-child dyad for a single item on the Diabetes Family Responsibility Questionnaire (DFRQ). This process was repeated for each of the 17 DFRQ items, and the individual item values were summed to arrive at the total agreement and discordance scores. Empty cells represent partial agreement between parent and child.

standardized against the Diabetes Control and Complications reference laboratory; the reference range is 4.0% to 6.0%.

Statistical Analysis

Statistical analyses were conducted using SAS (version 8.2 for Windows, SAS Institute, Inc., Cary, NC). Univariate analyses were examined first. An alpha level of .05 was used to determine statistical significance. The younger and older cohorts were compared using *t* tests. Pearson correlations were used to examine relationships between age, HbA1c, dyadic agreement, dyadic discordance, and diabetes-specific family conflict. A regression model was constructed to control for the complex relationships between variables, with the goal of further establishing the factors associated with glyce-mic control in this sample.

RESULTS

The mean age of the 121 youth was 12.1 ± 1.6 years; 50% were female. The study sample was ethnically and racially diverse, with 71% Caucasian, 12% African American, 10% Hispanic, and 7% reporting other or mixed ethnicity. The mean duration of diabetes was 5.4 ± 3.1 years, and the mean HbA1c was $8.4 \pm 1.4\%$. Pubertal status, by Tanner staging, was available for 73% of the sample. Of these 88 youth, 32% were prepubertal (Tanner I), 56% were pubertal (Tanner II–IV), and 13% were postpubertal (Tanner V). The mean ages of prepubertal, pubertal, and postpubertal youth were 10.7 ± 1.1 year, 12.0 ± 1.4 years, and 13.8 ± 0.6 years, respectively. The subset of youth with no Tanner staging data ($n = 33$) was significantly older than the 88 youth with Tanner staging data (12.9 ± 1.5 years vs. 11.8 ± 1.6 years, $p < .001$) and had a smaller proportion of girls (33% vs. 56%, $p < .03$).

To explore age-related differences in family behaviors between prepubertal and pubertal youth, we divided the sample into two groups. Because Tanner staging data

were only available for 73% of the sample and because of the differences in age and gender between those with and without Tanner staging data, we used age as a proxy for pubertal status when defining the two age groups. The younger cohort ($n = 57$) included youth younger than 12, and the older cohort ($n = 64$) included youth 12 and older. This division was consistent with the mean age of pubertal youth (12.0 ± 1.4 years). Indeed, when considering only the 88 youth with Tanner staging data, the younger cohort was 50% prepubertal and 50% pubertal, and the older cohort was 10% prepubertal, 62.5% pubertal, and 27.5% postpubertal, $\chi^2(2, N = 88) = 24.8, p < .0001$.

The mean age of the 57 youth in the younger cohort was 10.6 ± 0.7 years, and the mean age of the 64 youth in the older cohort was 13.5 ± 0.7 years. There were no significant differences between the two groups with respect to gender, duration of diabetes, or HbA1c. Table 2 displays characteristics of the entire sample and of the younger and older cohorts. Almost all of the participating parents or guardians were mothers ($n = 111, 91\%$); of the remaining adults, there were 8 fathers (7%), 1 stepmother (1%), and 1 grandmother (1%). A majority of parents or guardians had some education beyond high school: 13.5% had a graduate degree, 32% had a college degree, 35% had some college education, 17% had a high school diploma, and 2.5% had less than a high school diploma.

The dyadic variable of agreement, derived from the DFRQ, was not significantly correlated with age ($r = .15, p = .10$), nor did agreement differ significantly between age cohorts (see Table 3). However, discordance about sharing of responsibility was significantly inversely correlated with age ($r = -.24, p = .007$), with greater discordance associated with younger age, and there was significantly more discordance in the younger cohort than in the older cohort, $t(91) = 3.09, p = .003$ (see Table 3). Parent-reported diabetes-specific family conflict

Table 2

Means and Standards Deviations for Participant Characteristics for the Entire Sample and by Age Cohort

Characteristic	All participants (<i>N</i> = 121)	Younger cohort (<i>n</i> = 57)	Older cohort (<i>n</i> = 64)	<i>p</i> ^a
Gender (% female)	50	54	45	.32
Age (years)	12.1 ± 1.6	10.6 ± 0.7	13.5 ± 0.7	
Diabetes duration (years)	5.4 ± 3.1	5.2 ± 2.6	5.6 ± 3.6	.41
HbA1c (%)	8.4 ± 1.4	8.2 ± 1.4	8.7 ± 1.3	.08

Note. HbA1c = hemoglobin A1c.

^a Younger versus older cohort.

was positively correlated with youth age ($r = .21, p = .02$). However, when comparing the two age groups, although the mean conflict score was higher in the older cohort than in the younger cohort, the difference was not significant (see Table 3).

Pearson correlations indicated a significant relationship between parent report of diabetes-specific family conflict and glycemic control for the entire sample of families ($r = .36, p < .0001$), as well as for families in the younger cohort ($r = .28, p = .04$) and the older cohort ($r = .41, p = .0007$), with higher levels of reported conflict related to higher (worse) HbA1c levels in all groups (see Table 4). However, we did not find a similar pattern of relationships between youth report of diabetes-specific family conflict and HbA1c. The correlation between HbA1c and conflict as reported by youth in the younger cohort was not significant ($r = .15, p = .27$). Also, the correlation between HbA1c and conflict as reported by youth in the older cohort was not significant ($r =$

$.23, p = .06$). Another recent investigation has also reported a relationship between parental report of family behavior and adolescent HbA1c but not between adolescent report of family behavior and HbA1c (Cameron et al., 2008). Thus, we decided to focus only on parent report of conflict in the planned analyses by age group.

Dyadic agreement was significantly correlated with HbA1c levels in the entire sample ($r = -.21, p < .02$), with greater dyadic agreement related to lower (better) HbA1c levels. A breakdown by age cohort revealed that only in the younger cohort was dyadic agreement related to HbA1c levels ($r = -.32, p < .02$). Dyadic agreement and HbA1c levels were not significantly correlated for families in the older cohort. Dyadic discordance was not significantly correlated with HbA1c for the entire sample or for either age cohort. These relationships are reported in Table 4.

To further explore the relationship among dyadic agreement, diabetes-specific family conflict, and glycemic control in this

Table 3

Means and Standard Deviations for Total Parent–Child Dyadic Agreement and Discordance and Total Parent-Reported Diabetes-Specific Family Conflict for the Entire Sample and by Age Cohort

Family variables	All participants (<i>N</i> = 121)	Younger cohort (<i>n</i> = 57)	Older cohort (<i>n</i> = 64)	<i>p</i> ^a
Agreement about sharing of responsibility	9.3 ± 2.6	8.9 ± 2.9	9.7 ± 2.2	.08
Discordance about sharing of responsibility	0.77 ± 1.06	1.09 ± 1.25	0.50 ± 0.77	.003
Diabetes-specific family conflict (parent report)	6.4 ± 3.5	5.9 ± 3.5	6.8 ± 3.4	.16

^a Younger versus older cohort.

Table 4

Correlation of HbA1c With Dyadic Agreement, Dyadic Discordance, and Parent-Reported Diabetes-Specific Family Conflict for the Entire Sample and by Age Cohort

HbA1c with	All participants (<i>N</i> = 121)		Younger cohort (<i>n</i> = 57)		Older cohort (<i>n</i> = 64)	
	<i>r</i>	<i>p</i>	<i>r</i>	<i>p</i>	<i>r</i>	<i>p</i>
Dyadic agreement	-.21	.02	-.32	.02	-.16	<i>ns</i>
Dyadic discordance	.12	<i>ns</i>	.22	<i>ns</i>	.12	<i>ns</i>
Diabetes-specific conflict	.36	<.0001	.28	.04	.41	.0007

Note. HbA1c = hemoglobin A1c.

sample, we constructed a multivariate model with HbA1c as the dependent variable. Age group, gender, diabetes duration, parent report of diabetes-specific family conflict, and dyadic agreement were entered simultaneously in the model as independent variables. In a significant model, the independent variables accounted for 20% of the variance in HbA1c ($R^2 = .20$), $F(5) = 5.77$, $p < .0001$. Parent report of family conflict and dyadic agreement were the only independent predictors of HbA1c when controlling for potential confounders. Less family conflict ($p = .0003$) and greater dyadic agreement ($p = .02$) were associated with lower HbA1c (see Table 5).

DISCUSSION

The primary aim of this study was to explore the relationships between glycemic control, diabetes-specific family conflict, and the dyadic measures of agreement and discordance with respect to the sharing of diabetes management responsibilities in families with 9- to 14-year-old youth with

T1DM. In our sample from four clinical sites, there was significantly higher discordance for parent-child dyads in the younger, preteen cohort when compared with discordance in parent-child dyads in the older cohort. This suggests that abrupt changes in family sharing of diabetes management responsibilities are likely beginning to occur in the preteen age cohort. This finding is consistent with the qualitative research of Schilling et al. (2006), who also reported that parents begin to transfer responsibilities for diabetes management to children in late childhood, around 8 to 11 years of age.

Moreover, we found a relationship between agreement around responsibility sharing and glycemic control only for dyads in the younger cohort (9 to 11 years). In other words, only in the preteen cohort did we find a relationship between parent-child agreement with respect to responsibility sharing and glycemic control, with dyads with higher agreement having youth in the best glycemic control. It is possible

Table 5

*Summary of Regression Analysis for Variables Predicting Youth HbA1c (*N* = 121)*

Variable	B	SE B	β
Age group (younger vs. older)	0.41	0.23	0.15
Gender (female vs. male)	0.24	0.23	0.09
Diabetes duration	0.05	0.04	0.12
Diabetes-specific family conflict	0.12	0.03	0.32*
Dyadic agreement	-0.11	0.05	-0.20**

Note. Variables were entered simultaneously into the model, $F(5) = 5.77$, $R^2 = .20$, $p < .0001$. HbA1c = hemoglobin A1c.

* $p = .0003$. ** $p = .02$.

that agreement was significantly related to HbA1c only for this younger cohort because in the older cohort, glycemic control was more likely to be affected by pubertal hormones (Amiel et al., 1986). Supporting this conclusion is the significant difference in the proportion of pubertal or postpubertal youth in the two age cohorts. When considering only the 88 youth with available Tanner staging data, the younger cohort was 50% pubertal and 0% postpubertal, whereas the older cohort was 63% pubertal and 28% postpubertal.

In our entire sample, as well as within each age cohort, higher levels of parent-reported diabetes-specific family conflict were related to worse glycemic control in the youth. Other investigations have also found that higher levels of family conflict are related to worse glycemic control (Miller & Drotar, 2003). It has also been reported that family conflict with respect to diabetes management increases significantly from late childhood over the transition to adolescence (Anderson et al., 2002; Wysocki, Harris, Buckloh, Wilkinson, et al., 2006). In this sample of cross-sectional data, age was positively correlated with parent report of diabetes-specific family conflict. However, although parents in the older cohort reported more conflict than did parents in the younger cohort, the difference was not statistically significant.

Whereas dyadic measures of family behavior in diabetes can illuminate how key aspects of the parent-child relationship are changing over the transition to adolescence (Lerner et al., 1996), our study has several limitations. First, we derived dyadic variables from self-report data rather than from direct behavioral observations of interactions between parents and their 9- to 14-year-old children with T1DM. Moreover, our findings are based on cross-sectional data. Longitudinal studies are needed to prospectively track differences in dyadic measures in cohorts of older children transitioning to adolescence and their parents. Finally, some of the significant

correlations in our results are modest and await replication in other longitudinal or prospective studies. Despite these limitations, our study does identify diabetes-specific aspects of the parent-child relationship that are linked to glycemic outcomes and provides important information for future research in this area. Moreover, it is noteworthy that recent findings from a large European study of more than 2,000 adolescents and parents reported cross-sectional data very consistent with those of our study (Cameron et al., 2008).

Our findings have several clinical implications. Studying parent-child dyadic variables can help to identify potentially modifiable family variables, such as clear communication with respect to responsibility sharing for diabetes tasks that can be targeted in family-based interventions to optimize the glycemic control of youth in this vulnerable developmental period (Butler et al., 2008). Cameron et al. (2008) have observed that although many researchers have documented the relationship between family structure (e.g., number of parents in the home) and metabolic outcomes in adolescents with diabetes,

these aspects of family structure . . . are most intractable and least amenable to intervention by health professionals, and these investigators state that their finding of a relationship between discrepancies in parent and adolescent reports of responsibility for diabetes tasks “. . . demonstrated the importance of family dynamics in determining metabolic outcomes in Type 1 diabetes.” (p. 467)

Because most youth with T1DM are seen by a multidisciplinary team every 3 months for regular diabetes follow-up care (Silverstein et al., 2005), our findings suggest that clinicians engage older children and their parents in direct discussions about handling the transfer of responsibility for diabetes management tasks. Our findings, along with those of-

Cameron et al. (2008), suggest that pediatric diabetes clinicians should initiate discussions with transitioning youth and their parents to clarify who in the family is taking responsibility for the many different tasks involved in managing diabetes. The most recent Standards for the Care of Children and Adolescents With Type I Diabetes of the American Diabetes Association (Silverstein et al., 2005) recommended the following during the adolescent period: "continuing to involve parents appropriately with shared management . . . finding the degree of parental involvement that is comfortable for all involved, without risking deterioration in glycemic control from over- or underinvolvement" (p. 190). Our findings expand this recommendation to suggest that as youth with T1DM approach adolescence (at approximately 9 to 11 years old), the diabetes team should begin to engage parent-child dyads in discussions of how management responsibilities for diabetes tasks will be shared in the family and the optimal transfer of responsibilities over the early adolescent age period within each unique family system.

REFERENCES

- Allen, D. A., Tennen, H., McGrade, B. J., Afleck, G., & Ratzan, S. (1983). Parent and child perceptions of the management of juvenile diabetes. *Journal of Pediatric Psychology, 8*, 129–141.
- Amiel, S. A., Sherwin, R. S., Simonson, D. C., Lauritano, A. A., & Tamborlane, W. V. (1986). Impaired insulin action in puberty: A contributing factor to poor glycemic control in adolescents with diabetes. *New England Journal of Medicine, 315*, 215–219.
- Anderson, B. J., & Auslander, W. F. (1989). Research on diabetes management and the family: A critique. *Diabetes Care, 3*, 696–702.
- Anderson, B. J., Auslander, W. F., Jung, K. C., Miller, J. P., & Santiago, J. V. (1990). Assessing family sharing of diabetes responsibilities. *Journal of Pediatric Psychology, 15*, 477–492.
- Anderson, B. J., & Brackett, J. (2005). Diabetes in children. In F. J. Snoek & T. C. Skinner (Eds.), *Psychology in diabetes care* (2nd ed., pp. 1–25). West Sussex, England: Wiley.
- Anderson, B. J., Brackett, J., Ho, J., & Laffel, L. M. B. (1999). An office-based intervention to maintain parent-adolescent teamwork in diabetes management: Impact on parent involvement, family conflict, and subsequent glycemic control. *Diabetes Care, 22*, 713–721.
- Anderson, B. J., Ho, J., Brackett, J., Finkelstein, D., & Laffel, L. (1997). Parental involvement in diabetes management tasks: Relationships to blood glucose monitoring adherence and metabolic control in young adolescents with insulin-dependent diabetes mellitus. *Journal of Pediatrics, 130*, 257–265.
- Anderson, B. J., Vangsness, L., Connell, A., Butler, D., Goebel-Fabbri, A., & Laffel, L. M. B. (2002). Family conflict, adherence, and glycemic control in youth with short duration Type 1 diabetes. *Diabetic Medicine, 19*, 635–642.
- Barber, B. K. (1996). Parental psychological control: Revisiting a neglected construct. *Child Development, 67*, 3296–3319.
- Beck, J. S., Beck, A. T., & Jolly, J. (2001). Manual for the Beck Youth Inventories of Emotional and Social Adjustment. San Antonio, TX: Psychological Corporation.
- Butler, D. A., Zuehlke, J. B., Tovar, A., Volkening, L. K., Anderson, B. J., & Laffel, L. M. B. (2008). The impact of modifiable family factors on glycemic control among youth with Type 1 diabetes. *Pediatric Diabetes, 9*, 373–381.
- Cameron, F. J., Skinner, T. C., de Beaufort, C. E., Hoey, H., Swift, P. G., Aanstoot, H., et al. (2008). Are family factors universally related to metabolic outcomes in adolescents with Type 1 diabetes? *Diabetic Medicine, 5*, 463–468.
- Chase, H. P. (2006). *Understanding diabetes* (11th ed.). Denver, CO: Children's Diabetes Foundation at Denver.
- Cook, W., & Kenny, D. A. (2006). Examining the validity of self-report assessment of family functioning: A question of the level of analysis. *Journal of Family Psychology, 20*, 209–216.
- Cox, D., Irvine, A., Gonder-Frederick, L., Nowacek, G., & Butterfield, J. (1987). Fear of hypoglycemia: Quantification, validation, and utilization. *Diabetes Care, 10*, 617–621.
- Diabetes Control and Complications Trial Research Group. (1993). The effect of intensive treatment of diabetes on the development and progression of long-term complications in insulin-dependent diabetes mellitus. *New England Journal of Medicine, 329*, 977–986.
- Ellis, D. A., Frey, M., Narr-King, S., Templin, T., Cunningham, P. B., & Cakan, N. (2005).

- The effects of multisystemic therapy on diabetes stress among adolescents with chronically poorly controlled Type 1 diabetes: Findings from a randomized, controlled trial. *Pediatrics*, *116*, 826–832.
- Epidemiology of Diabetes Interventions and Complications Research Group. (2001). Beneficial effect of intensive therapy of diabetes during adolescence: Outcomes after the conclusion of the Diabetes Control and Complications Trial (DCCT). *Journal of Pediatrics*, *139*, 804–812.
- Harris, M. A., Wysocki, T., Sadler, M., Wilkinson, K., Harvey, L. M., Buckloh, L. M., et al. (2000). Validation of a structured interview for the assessment of diabetes self-management. *Diabetes Care*, *23*, 1301–1304.
- Hartos, J. L., Eitel, P., & Simons-Morton, B. (2001). Do parent-imposed delayed licensure and restricted driving reduce risky driving behaviors among newly licensed teens? *Prevention Science*, *2*, 113–122.
- Hauser, S. T., Jacobson, A. M., Lavori, P., Wolfson, J. I., Herskowitz, R. D., Milley, J., et al. (1990). Adherence among children and adolescents with insulin-dependent diabetes mellitus over a four-year longitudinal follow-up: II. Immediate and long-term linkages with the family milieu. *Journal of Pediatric Psychology*, *15*, 527–542.
- Holmbeck, G. N. (1996). A model of family relational transformations during the transition to adolescence: Parent-adolescent conflict and adaptation. In J. A. Graber, J. Brooks-Gunn, & A. C. Petersen (Eds.), *Transitions through adolescence: Interpersonal domains and context* (pp. 167–199). Mahwah, NJ: Erlbaum.
- Hood, K. K., Butler, D. A., Anderson, B. J., & Laffel, L. M. B. (2007). Updated and revised Diabetes Family Conflict Scale. *Diabetes Care*, *30*, 1764–1769.
- Hood, K. K., Butler, D. A., Volkening, L. K., Anderson, B. J., & Laffel, L. M. B. (2004). The Blood Glucose Monitoring Communication Questionnaire: An instrument to measure affect specific to blood glucose monitoring. *Diabetes Care*, *27*, 2610–2615.
- Iannotti, R. J., Schneider, S., Nansel, T. R., Haynie, D. L., Plotnick, L. P., Clark, L. M., et al. (2006). Self-efficacy, outcome expectations, and diabetes self-management in adolescents with Type 1 diabetes. *Journal of Developmental and Behavioral Pediatrics*, *27*, 98–105.
- Ingersoll, G. M., Orr, D. P., Herrold, A. J., & Golden, M. P. (1986). Cognitive maturity and self-management among adolescents with insulin-dependent diabetes mellitus. *Journal of Pediatrics*, *108*, 620–623.
- Jackson, C., Henriksen, L., & Foshee, V. A. (1998). The Authoritative Parenting Index: Predicting health risk behaviors among children and adolescents. *Health Education & Behavior*, *25*, 319–337.
- Johnson, S. B., Silverstein, J., Rosenbloom, A., Carter, R., & Cunningham, W. (1986). Assessing daily management in childhood diabetes. *Health Psychology*, *5*, 545–564.
- Kovacs, M. (1985). The Children's Depression Inventory (CDI). *Psychopharmacology Bulletin*, *21*, 995–998.
- Lerner, R. M., Lerner, J. V., von Eye, A., Ostrum, C. W., Nitz, K., Talwar-Soni, R., et al. (1996). Continuity and discontinuity across the transition of early adolescence: A developmental contextual perspective. In J. A. Graber, J. Brooks-Gunn, & A. C. Petersen (Eds.), *Transitions through adolescence: Interpersonal domains and context* (pp. 3–22). Mahwah, NJ: Erlbaum.
- Levitt, M. J., Guacci-Franco, N., & Levitt, J. L. (1993). Convoys of social support in childhood and early adolescence: Structure and function. *Developmental Psychology*, *29*, 811–818.
- Matheny, A., Wachs, T., Ludwig, J., & Phillips, K. (1995). Bringing order out of chaos: Psychometric characteristics of the Confusion, Hubbub, and Order Scale. *Journal of Applied Developmental Psychology*, *16*, 429–444.
- Miller, V. A., & Drotar, D. (2003). Discrepancies between mother and adolescent perceptions of diabetes-related decision-making autonomy and their relationship to diabetes-related conflict and adherence to treatment. *Journal of Pediatric Psychology*, *28*, 265–274.
- Nansel, T. R., Anderson, B. J., Laffel, L. M. B., Simons-Morton, B. G., Weissberg-Benchell, J., Wysocki, T., et al. (2009). A multisite trial of a clinic-integrated intervention for promoting family management of pediatric Type 1 diabetes: Feasibility and design. *Pediatric Diabetes*, *10*, 105–115. doi: 10.1111/j.1399-5448.2008.00448.x.
- Nansel, T. R., Rovner, A. J., Haynie, D., Iannotti, R. J., Simons-Morton, B., Wysocki, T., et al. (2009). Development and validation of the Collaborative Parent Involvement scale for youths with Type 1 diabetes. *Journal of Pediatric Psychology*, *34*, 30–40. doi: 10.1093/jpepsy/jsn058.
- Palmer, D. L., Berg, C. A., Wiebe, D. J., Beveridge, R. M., Korb, C. D., Upchurch, R., et al. (2004). The role of autonomy and pubertal

- status in understanding age differences in maternal involvement in diabetes responsibility across adolescence. *Journal of Pediatric Psychology*, *29*, 35–46.
- Piazza-Waggoner, C., Modi, A. C., Powers, S. W., Williams, L. B., Dolan, L. M., & Patton, S. R. (2008). Observational assessment of family functioning in families with children who have Type 1 diabetes mellitus. *Journal of Developmental and Behavioral Pediatrics*, *29*, 101–105.
- Rubin, R., Young-Hyman, D., & Peyrot, M. (1989). Parent-child responsibility and conflict in diabetes care. *Diabetes*, *38*(Suppl. 2), 28.
- Schilling, L. S., Knafelz, K. A., & Grey, M. (2006). Changing patterns of self-management in youth with Type 1 diabetes. *Journal of Pediatric Nursing*, *21*, 412–424.
- Silverstein, J., Klingensmith, G., Copeland, K., Plotnick, L., Kaufman, F., Laffel, L., et al. (2005). Care of children and adolescents with Type 1 diabetes: A statement of the American Diabetes Association. *Diabetes Care*, *28*, 186–212.
- Skinner, T. C., Murphy, H., & Huws-Thomas, M. (2005). Diabetes in adolescents. In F. J. Snoek & T. C. Skinner (Eds.), *Psychology in diabetes care* (2nd ed., pp. 27–52). West Sussex, England: Wiley.
- Smetana, J., & Asquith, P. (1994). Adolescents' and parents' conceptions of parental authority and personal autonomy. *Child Development*, *65*, 1147–1162.
- Varni, J., Burwinkle, T., Jacobs, J., Gottschalk, M., Kaufman, F., & Jones, K. (2003). The PedsQL in type 1 and type 2 diabetes. *Diabetes Care*, *26*, 631–637.
- Varni, J. W., Seid, M., & Kurtin, P. S. (2001). The PedsQL™ 4.0: Reliability and validity of the Pediatric Quality of Life Inventory™ Version 4.0 Generic Core Scales in healthy and patient populations. *Medical Care*, *39*, 800–812.
- Wiebe, D. J., Berg, C. A., Korb, C., Palmer, D. L., Beveridge, R. M., Upchurch, R., et al. (2005). Children's appraisals of maternal involvement in coping with diabetes: Enhancing our understanding of adherence, metabolic control, and quality of life across adolescence. *Journal of Pediatric Psychology*, *30*, 167–178.
- Wysocki, T., Harris, M. A., Buckloh, L. M., Mertlich, D., Lochrie, A. S., Taylor, A., et al. (2006). Effects of behavioral family systems therapy for diabetes on adolescents' family relationships, treatment adherence, and metabolic control. *Journal of Pediatric Psychology*, *31*, 928–938.
- Wysocki, T., Harris, M. A., Buckloh, L. M., Wilkinson, K., Sadler, M., Mauras, N., et al. (2006). Self-care autonomy and outcomes of intensive therapy or usual care in youth with Type 1 diabetes. *Journal of Pediatric Psychology*, *31*, 1036–1045.
- Wysocki, T., Taylor, A., Hough, B. S., Linscheid, T. R., Yeates, K. O., & Naglieri, J. A. (1996). Deviation from developmentally appropriate self-care autonomy: Association with diabetes outcomes. *Diabetes Care*, *19*, 119–125.

E-Mail Notification of Your Latest Issue Online!

Would you like to know when the next issue of your favorite APA journal will be available online? This service is now available to you. Sign up at <http://notify.apa.org/> and you will be notified by e-mail when issues of interest to you become available!